



Problem 1 Define A = event exactly 9 people say yes.

1. The complement of A is the event that:
 - a) anything other than 9 people say yes
 - b) anything other than 9 people say no
 - c) exactly 6 people say yes
 - d) exactly 6 people say no
 - e) more than 6 people say no

Problem 2 At a large university, the probability that a student takes calculus and statistics in the same semester is 0.0125. The probability that a student takes statistics is 0.125; the probability that a student takes calculus is 0.3

2. Find the probability that a student is taking calculus, given that he or she is taking statistics.
 - a) 0.1
 - b) 0.1125
 - c) 0.0016
 - d) 0.1375
 - e) 0.4800
3. Is the event of taking calculus independent of the event of taking statistics? Justify your answer numerically.

Problem 3

- 11) If you flip a coin three times, the possible outcomes are HHH HHT HTH HTT THH THT TTH TTT. What is the probability of getting at least one head?
- A) $\frac{1}{2}$ B) $\frac{7}{8}$ C) $\frac{1}{4}$ D) $\frac{3}{4}$

- 12) When a quarter is tossed four times, 16 outcomes are possible.

HHHH	HHHT	HHTH	HHTT
HTHH	HTHT	HTTH	HTTT
THHH	THTT	THTH	THTT
TTHH	TTHT	TTTH	TTTT

Here, for example, HHTH represents the outcome that the first toss is heads, the next two tosses are tails, and the fourth toss is heads. The events A and B are defined as follows.

Event A = the first two tosses are heads
Event B = the first and last tosses are the same

Are the events A and B mutually exclusive?

- A) Yes B) No



Problem 4

The American Community Survey is an ongoing survey that provides data every year to give communities the current information they need to plan investments and services. The 2010 American Community Survey estimates that 14.6% of Americans live below the poverty line, 20.7% speak a language other than English (foreign language) at home, and 4.2% fall into both categories.

4. Are living below the poverty line and speaking a foreign language at home disjoint?
5. What percent of Americans live below the poverty line and only speak English at home?
6. What percent of Americans live below the poverty line or speak a foreign language at home?
7. What percent of Americans live above the poverty line and only speak English at home?
8. Is the event that someone lives below the poverty line independent of the event that the person speaks a foreign language at home?

Problem 5

Consider $P(A) = 0.3$, $P(B) = 0.7$

9. Can you compute $P(A \text{ and } B)$ if you only know $P(A)$ and $P(B)$?
10. Assuming that events A and B arise from independent random processes,
 - a. What is $P(A \text{ and } B)$?
 - b. What is $P(A \text{ or } B)$?
 - c. What is $P(A|B)$?
 - d. If we are given that $P(A \text{ and } B) = 0.1$, are the random variables giving rise to events A and B independent?
 - e. If we are given that $P(A \text{ and } B) = 0.1$, what is $P(A|B)$?

Problem 6

A quiz consists of 3 multiple choice questions with 4 possible answers each. If a student guesses the answer to each question, what is the probability that they get all the answers correct?

Problem 7¹

Topics: contingency table, conditional probability, joint probability, marginal probability.

The family college data set contains a sample of 792 cases with two variables, teen and parents, and is summarized in the following Table:

		parents		Total
		degree	not	
teen	college	231	214	445
	not	49	298	347
Total		280	512	792

¹ Open Intro Stat Book



The teen variable is either college or not, where the college label means the teen went to college immediately after high school. The parent's variable takes the value degree if at least one parent of the teenager completed a college degree.

11. If at least one parent of a teenager completed a college degree, what is the chance the teenager attended college right after high school?
12. Probability of a random teenager from the study went to college:

Problem 8¹

Topics: general multiplication rule

A smallpox data set provides a sample of 6,224 individuals from the year 1721 who were exposed to smallpox in Boston. Doctors at the time believed that inoculation, which involves exposing a person to the disease in a controlled form, could reduce the likelihood of death.

Suppose we are given only two pieces of information: 96.08% of residents were not inoculated, and 85.88% of the residents who were not inoculated ended up surviving.

13. How could we compute the probability that a resident was not inoculated and lived?

¹ Open Intro Stat Book