



# Week in Review

## Math 152

---

### **Week 03**

Volumes by Slicing: Disks and Washers  
Volume by Cylindrical Shells



# General slicing method for any volume

**Step 1:** Plot the graph

**Step 2.** Find the size of a slice at  $x$  or  $y$

- At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$
- At  $y$ : Thickness  $dy \Rightarrow$  Cross section  $A(y)$

**Step 3.** Find the volume of a slice at  $x$  or  $y$



- At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$ 
  - $dV = A(x)dx$
- At  $y$ : Thickness  $dy \Rightarrow$  Cross section  $A(y)$ 
  - $dV = A(y)dy$

**Step 4.** Find the upper/lower limits for  $x$  or  $y$

**Step 5.** Set up integral and evaluate

$$V = \int_a^b A(x)dx \text{ or } V = \int_c^d A(y)dy$$

The solid whose base is the region bounded by semicircle  $y = \sqrt{1 - x^2}$  and the  $x$ -axis. And whose cross section through the solid perpendicular to the  $x$  axis are squares. Find the volume of the solid.



$$A(x) = [\sqrt{1 - x^2}]^2$$

$$V(\text{slice}) = A(x)dx = (1 - x^2)dx$$

- Limit for  $x$  :  $[-1,1]$
- $V = \int_a^b A(x)dx$ 


$$= \int_{-1}^1 (1 - x^2)dx = 2 \int_0^1 (1 - x^2)dx$$

$$= 2 \left[ x - \frac{1}{2x^2} \right]_0^1 = 1$$



## Example

Find the volume of the solid whose base is the ellipse  $x^2 + 4y^2 = 4$  and whose cross-sections perpendicular to the  $y$ -axis are squares. Evaluate your integral.





## Example

The base of a solid is the region bounded by the curve  $y = 5 - x^2$  and the  $x$ -axis. Cross-Sections perpendicular to the  $y$ -axis are rectangles with height equal to twice the base. Find the volume of this solid.



## Example

Consider the solid  $S$  whose base is the region bounded by  $y = 4 - x^2$  and  $y = 0$ . Cross sections perpendicular to the  $y$  - axis are semicircles. Find the volume of  $S$ .



## Example

Consider the solid  $S$  described here. The base of  $S$  is the region bounded by  $y = x^2$  and  $y = 4$ . Cross sections perpendicular to the  $x$  - axis are squares. Find the volume of  $S$ .



# Volume by disks perpendicular to the x axis

## Volume of solid of revolution around **x axis**

**Step 1:** Plot the graph

**Step2.** Find the size of a perpendicular slice at  $x$

At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$

$$\pi[f(x)]^2$$

**Step3.** Find the volume of a slice at  $x$

• At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$


- $dV = \pi[f(x)]^2 dx$

**Step4.** Find the upper/lower limits for  $x$

**Step5.** Set up integral and evaluate

$$V = \int_a^b A(x)dx = \int_a^b \pi[f(x)]^2 dx$$

Find the volume of the solid that is obtained when the region under the curve  $y = \sqrt{x}$  over the interval  $[1, 4]$  is revolved about the x-axis



$$V(\text{disk}) = \pi[f(x)]^2 dx$$

$$1 \leq x \leq 4 \text{ (limits)}$$

$$\begin{aligned} V &= \int_a^b \pi[f(x)]^2 dx \\ &= \int_1^4 \pi x dx \\ &= \frac{\pi}{2} [x^2]_1^4 \\ &= \frac{\pi}{2} [16 - 1] = \frac{15}{2} \pi \end{aligned}$$



# Volume by Washer perpendicular to the x axis

**Volume of solid of revolution around  $x$  axis**

**Step 1:** Plot the graph of  $f(x)$ ,  $g(x)$  w/  $f > g$

**Step 2.** Find the size of a perpendicular slice at  $x$

- At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$
- Washer = Large disc – small disc  

$$= \pi[f(x)]^2 - \pi[g(x)]^2$$

**Step 3.** Find the volume of a slice at  $x$

- At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(x)$ 
  - $dV = \pi([f(x)]^2 - [g(x)]^2)dx$



**Step 4.** Find the upper/lower limits for  $x$

**Step 5.** Set up integral and evaluate

$$V = \int_a^b A(x)dx$$

$$= \int_a^b \pi([f(x)]^2 - [g(x)]^2)dx$$

Find the volume of the solid that is obtained when the region between the curve  $y = \sqrt{x}$  and  $y = 2$  over the interval  $[1, 4]$  is revolved about the  $x$ -axis



$$V(\text{washer}) = \pi\{\sqrt{x}\}^2 dx - \pi\{1\}^2 dx = \pi[x - 1]dx$$

Limit =  $[1, 4]$

$$V = \int_1^4 \pi(x - 1)dx$$

$$= \pi \left[ \frac{1}{2}x^2 - x \right]_1^4 = \frac{9}{2}\pi$$





# Example

The region bounded by  $y = \cos x$  and the  $x$ -axis on the interval  $\left[0, \frac{\pi}{2}\right]$  is rotated about the  $x$ -axis. Find the volume of the resulting solid.

- (a) 1
- (b)  $\frac{\pi^2}{2}$
- (c)  $\frac{\pi}{2}$
- (d)  $\frac{\pi}{4}$
- (e)  $\frac{\pi^2}{4}$  ← correct



## Example

The region bounded by  $y = e^x$  and the  $x$ -axis on the interval  $[0, 2]$  is rotated about the  $x$ -axis. Find the volume of the resulting solid.

(a)  $\frac{\pi e^4}{2}$

(b)  $\frac{\pi e^2}{2}$

(c)  $\frac{\pi}{2}(e^4 - 1)$  ← correct

(d)  $\frac{\pi}{2}(e^2 - 1)$

(e)  $2\pi(e^4 - 1)$



## Example

The region bounded by the curves  $y = x^2$  and  $y = 1$  is rotated about the line  $y = 1$ . Find the volume of the resulting solid.

- (a)  $\frac{8\pi}{15}$
- (b)  $\frac{8\pi}{5}$
- (c)  $\frac{4\pi}{3}$
- (d)  $\frac{12\pi}{5}$
- (e)  $\frac{16\pi}{15}$



## Example

If we revolve the region bounded by  $y = 1 - x^2$  and  $x - y = 1$  about the line  $y = 3$ , which of the following integrals gives the resulting volume?

(a)  $\int_{-1}^2 2\pi(3-x)(x^2-x+2) dx$

(b)  $\int_{-2}^1 \pi((2+x^2)^2 - (4-x)^2) dx$

(c)  $\int_{-1}^2 2\pi(x-3)(x^2-x+2) dx$

(d)  $\int_{-2}^1 \pi((4-x)^2 - (2+x^2)^2) dx$

(e)  $\int_{-1}^2 \pi((2+x^2)^2 - (4-x)^2) dx$



## Example

Consider the region  $R$  bounded by  $y = \sqrt{x}$ ,  $y = 1$ ,  $x = 0$ . Find the volume obtained by rotating the region  $R$  about the line  $y = 1$ .

- (a)  $\frac{\pi}{6}$
- (b)  $\frac{\pi}{2}$
- (c)  $\frac{7\pi}{6}$
- (d)  $\frac{\pi}{3}$
- (e)  $\frac{5\pi}{6}$



# Example

Which of the following integrals gives the volume of the solid obtained by rotating the region bounded by  $y = 5 - x^2$  and  $y = 1$  about the  $x$ -axis.

(a)  $\pi \int_{-2}^2 (1 - (5 - x^2)^2) dx$

(b)  $\pi \int_{-2}^2 (4 - x^2)^2 dx$

(c)  $2\pi \int_{-2}^2 x(4 - x^2) dx$

(d)  $\pi \int_{-2}^2 ((5 - x^2)^2 - 1) dx$  ← correct

(e)  $2\pi \int_{-2}^2 x(x^2 - 4) dx$



# Volume by disks perpendicular to the y axis

**Volume of solid of revolution around y axis**

**Step 1:** Plot the graph

**Step 2.** Find the size of a perpendicular slice at y

- At y: Thickness  $dy \Rightarrow$  Cross section  $A(y)$   
 $\pi[g(y)]^2$
- For  $y = f(x)$ , solve for  $x = f^{-1}(y)$

**Step 3.** Find the volume of a slice at y


- At y: Thickness  $dy \Rightarrow$  Cross section  $A(y)$ 
  - $dV = \pi[g(y)]^2 dy$

**Step 4.** Find the upper/lower limits for y

**Step 5.** Set up integral and evaluate

$$V = \int_c^d A(y) dy = \int_c^d \pi[g(y)]^2 dy$$

Find the volume of the solid that is obtained when the region between the curve  $y = x^3$  and  $x = 0$  between the interval  $0 \leq y \leq 8$  is revolved about the y-axis



$$A(x) = \pi(\sqrt[3]{y})^2$$

Radius =  $\sqrt[3]{y}$

$$V(\text{---}) = \pi[g(y)]^2 dy$$

$$0 \leq y \leq 8 \text{ (limits)}$$

$$\begin{aligned} V &= \int_a^b \pi[g(y)]^2 dy \\ &= \int_0^8 \pi y^{\frac{2}{3}} dy \\ &= \pi \left[ \frac{3}{5} x^{\frac{5}{3}} \right]_0^8 \end{aligned}$$



# Volume by Washer perpendicular to the y axis

**Volume of solid of revolution around y axis**

**Step 1:** Plot the graph of  $f(y)$ ,  $g(y)$  w/  $f > g$

**Step 2.** Find the size of a perpendicular slice at  $y$

- At  $y$ : Thickness  $dy \Rightarrow$  Cross section  $A(y)$
- Washer = Large disc – small disc  

$$= \pi[f(y)]^2 - \pi[g(y)]^2$$

**Step 3.** Find the volume of a slice at  $y$

- At  $x$ : Thickness  $dx \Rightarrow$  Cross section  $A(y)$ 
  - $dV = \pi([f(y)]^2 - [g(y)]^2)dy$




**Step 4.** Find the upper/lower limits for  $y$

**Step 5.** Set up integral and evaluate

$$V = \int_a^b A(y)dy$$

$$= \int_a^b \pi([f(y)]^2 - [g(y)]^2)dy$$

Find the volume of the solid that is obtained when the region between the curve  $y = x^2$  and  $y = 2x$  is revolved about the y-axis



||

$$A(x) = \pi\{\sqrt{y}\}^2 - \pi\{y/2\}^2 dy$$

$$A(y) = \pi\{1/2 y\}^2 - \pi\{y\}^2 dy$$

$$V(\text{washer}) = \pi\{\sqrt{x}\}^2 - \pi\{1\}^1 = \pi[x - 1]$$

$$\text{Limit} = [1,4]$$

$$V = \int_1^4 \pi(x - 1)dx$$

$$= \pi \left[ \frac{1}{2}x^2 - x \right]_1^4 = \frac{9}{2}\pi$$





# Example

Find the volume of the solid obtained by rotating the region bounded by  $x = y^2$  and  $x = y^3$  around the  $y$ -axis.

- (a)  $\frac{\pi}{35}$
- (b)  $\frac{\pi}{10}$
- (c)  $\frac{\pi}{12}$
- (d)  $\frac{2\pi}{35}$  ← correct
- (e)  $\frac{\pi}{105}$



## Example

Consider the region  $R$  bounded by  $y = 2x^2$  and  $y = 1$ , first quadrant only.

Find the volume obtained by rotating  $R$  about the  $y$ -axis.

- (a)  $\frac{\pi}{4}$
- (b)  $\frac{\pi}{2}$
- (c)  $\pi$
- (d)  $\frac{4\pi}{5}$
- (e) None of the above



# Volume by cylindrical shells about the x-axis

**Volume of solid of revolution around x axis**

**Step 1:** Plot the graph

**Step 2.** Find the size of a parallel slice at  $y$

At  $y$ : Thickness  $dy \Rightarrow$  Cross section  $A(y)$

$$2\pi f(y)dy$$

**Step 3.** Find the volume of a slice at  $y$

• At  $y$ : Thickness  $dx \Rightarrow$  Cross section  $A(y)$


- $dV = 2\pi f(y)dy$

**Step 4.** Find the upper/lower limits for  $y$

**Step 5.** Set up integral and evaluate

$$V = \int_a^b A(y)dy = \int_a^b 2\pi y f(y)dy$$

Use cylindrical shells to find the volume of the solid generated when the shaded region is revolved about the indicated axis.



$$\begin{aligned}
 V &= \int_0^2 2\pi y[y - y^2 + 2]dy \\
 &= 2\pi \int_0^2 [y^2 - x^3 + 2y]dy \\
 &= 2\pi \left[ \frac{1}{3}y^3 - \frac{1}{4}y^4 + y^2 \right]_0^2 \\
 &= 2\pi \left[ \frac{8}{3} - \frac{16}{4} + 4 \right] \\
 &= \frac{16\pi}{3}
 \end{aligned}$$



## Example

Consider the region bounded by the two curves  $y = \cos x$ ,  $y = \sin x$  and the two lines  $x = 0$  and  $x = \frac{\pi}{4}$ . Which of the following represents the volume of this region being rotated about the line  $x = -1$ ?

(a)  $\int_0^{\frac{\pi}{4}} 2\pi(x+1)(\cos x - \sin x) dx$  ← correct

(b)  $\int_0^{\frac{\pi}{4}} 2\pi(x+1)(\sin x - \cos x) dx$

(c)  $\int_{-1}^{\frac{\pi}{4}} 2\pi(x+1)(\cos x - \sin x) dx$

(d)  $\int_0^{\frac{\pi}{4}} 2\pi(x+1)(\cos^2 x - \sin^2 x) dx$

(e)  $\int_0^{\frac{\pi}{4}} \pi(\cos^2 x - \sin^2 x) dx$



## Example

Find the volume of the solid found by rotating the region bounded by the curves  $y = -x^2 + 2x$  and  $y = 0$  about the  $y$ -axis.

- (a)  $\frac{16}{3}\pi$
- (b)  $\frac{8}{3}\pi$
- (c)  $\frac{4}{3}\pi$
- (d)  $\frac{2}{3}\pi$
- (e)  $\frac{1}{3}\pi$



## Example

Consider the region  $R$  bounded by  $y = x^3$ ,  $y = -x + 2$ ,  $x = 0$ , and  $x = 1$ .

- (a) Sketch the region  $R$ .
- (b) Set up the integral that gives the volume obtained by revolving the region  $R$  about the  $x$ -axis using the method of washers. **DO NOT EVALUATE THE INTEGRAL.**
- (c) Set up the integral that gives the volume obtained by revolving the region  $R$  about the line  $x = 1$  using the method of cylindrical shells. **DO NOT EVALUATE THE INTEGRAL.**



## Example

Consider the region  $R$  bounded by  $u = 4x - x^2$  and  $u = 0$ . Which of the following integrals gives the volume of the solid obtained by revolving  $R$  about the line  $x = -2$ ?

(a)  $\int_0^4 2\pi(2-x)(4x-x^2) dx$

(b)  $\int_0^4 2\pi x(4x-x^2) dx$

(c)  $\int_0^4 2\pi(x+2)(4x-x^2) dx$

(d)  $\int_0^4 2\pi(x-2)(4x-x^2) dx$

(e) None of the above



## Example

Consider the region  $R$  bounded by  $y = \ln x$ ,  $y = 0$ , and  $x = 2$ . If this region is revolved about the line  $y = -2$ :

- (a) Set up but **do not evaluate** the integral that gives the volume using the method of shells.
- (b) Set up but **do not evaluate** the integral that gives the volume using the method of washers.





## Example

Consider the region bounded by the curves  $x = y^2 - 2y$  and the  $y$ -axis. Which of the following represents the volume of solid formed when the region is rotated about  $y = 4$ ?

(a)  $\int_0^2 2\pi y(y^2 - 2y) dy$

(b)  $\int_0^2 2\pi y(2y - y^2) dy$

(c)  $\int_0^2 2\pi(4 - y)(y^2 - 2y) dy$

(d)  $\int_0^2 \pi(y - 4)(4y^2 - y^4) dy$

(e)  $\int_0^2 2\pi(4 - y)(2y - y^2) dy$  ← correct