



MATH 140: WEEK-IN-REVIEW 10 (CHAPTERS 5.5 & 5.6)

1. Compute the following values for the function $f(x) = \begin{cases} 5x - 3 & \text{if } x < -4, \\ 2x^2 - 1 & \text{if } -4 \leq x \leq 4, \\ 6 & \text{if } 4 < x \leq 6, \\ \frac{3}{x-4} & \text{if } x > 6. \end{cases}$

(a) $f(-5) = 5(-5) - 3$
 $= -25 - 3$
 $= -28$

(b) $f(-4) = 2(-4)^2 - 1 = 2 \cdot 16 - 1$
 $= 32 - 1$
 $= 31$

(c) $f(0) = 2(0)^2 - 1$
 $= -1$

(d) $f(4) = 2(4)^2 - 1$
 $= 2 \cdot 16 - 1$
 $= 32 - 1 = 31$

(e) $f(8)$ DNE since $x=8$ is not in domain

(f) $f(9) = \frac{3}{9-4} = \frac{3}{5}$



2. State the domain of the function $g(x) = \begin{cases} 5x+7 & \text{if } x < -4, \text{ A} \\ \frac{1}{3x+7} & \text{if } -3 \leq x < 3, \text{ B} \\ -5\sqrt{x-2} & \text{if } x \geq 3. \text{ C} \end{cases}$

In A, $5x+7$ is a polynomial \Rightarrow no restrictions : $(-\infty, -4)$

In B, $\frac{1}{3x+7}$ is a rational function \Rightarrow denominator $\neq 0$, $3x+7 \neq 0 \Rightarrow \frac{3x}{3} \neq \frac{-7}{3}$
 $x \neq -\frac{7}{3}$: $[-3, -\frac{7}{3}) \cup (-\frac{7}{3}, 3)$


In C, $-5\sqrt{x-2}$ is an even root function $\Rightarrow x-2 \geq 0 \Rightarrow x \geq 2$. But in C, $x \geq 3$
 : $[3, \infty)$

Domain of $g(x)$: $(-\infty, -4) \cup [-3, \frac{7}{3}) \cup (-\frac{7}{3}, \infty)$

3. Sketch the graph of $h(x) = \begin{cases} -2x-3 & \text{if } x \leq -3, \text{ A} \\ -x^2-4x+5 & \text{if } -3 < x < 1, \text{ B} \rightarrow \text{vertex } (-2, 9) \\ 6 & \text{if } x \geq 2. \text{ C} \end{cases}$

A: $-2x-3$ (line)

x	f(x)	(x,y)
-3	$-2(-3)-3 = 3$	(-3, 3)
-4	$-2(-4)-3 = 5$	(-4, 5)
-5	$-2(-5)-3 = 7$	(-5, 7)



B: $-x^2-4x+5$ (Quadratic fn)

x	f(x)	(x,y)
-3	$-(-3)^2-4(-3)+5 = -9+12+5 = 8$	(-3, 8) hole
-2	$-(-2)^2-4(-2)+5 = -4+8+5 = 9$	(-2, 9)
0	$-0^2-4\cdot 0+5 = 5$	(0, 5)
1	$-1^2-4\cdot 1+5 = 0$	(1, 0) hole




4. Rewrite the function $f(x) = |18 - 3x|$ as a piecewise-defined function.

$$\begin{aligned}
 |18 - 3x| &= \begin{cases} -1 \cdot (18 - 3x) & \text{if } 18 - 3x < 0 \\ 18 - 3x & \text{if } 18 - 3x \geq 0 \end{cases} \\
 &= \begin{cases} -18 + 3x & \text{if } x > 6 \\ 18 - 3x & \text{if } x \leq 6 \end{cases} \\
 &= \begin{cases} 18 - 3x & \text{if } x \leq 6 \\ -18 + 3x & \text{if } x > 6 \end{cases} \quad * \text{ in order of interval}
 \end{aligned}$$

$18 - 3x < 0$
 $\frac{18}{3} < \frac{3x}{3}$
 $6 < x$

5. A truck rental agency charges a flat fee of \$20. If the distance traveled is less than 100 miles, the cost is \$0.95 per mile. For any distance greater than 100 miles, the cost reduces to \$0.85 per each additional mile. Write a function $C(m)$ representing the cost, C , for m miles driven.



$$C(m) = \begin{cases} 20 + 0.95m & \text{if } 0 \leq m \leq 100 \\ 115 + 0.85(m - 100) & \text{if } m > 100 \end{cases}$$



6. Rewrite each exponential expression as a single equivalent expression in the given base.

(a) $7 \cdot 49^{x+2}$ base 7. $49 = 7^2$

$$7 \cdot (7^2)^{x+2}$$

$$= 7^1 \cdot 7^{2(x+2)} = 7^1 \cdot 7^{2x+4} = 7^{2x+5}$$

(b) $\left(\frac{1}{3}\right)^x \cdot \frac{27}{9^x}$ base 3. $27 = 3^3$, $9 = 3^2$

$$\left(\frac{1}{3}\right)^x \cdot \frac{3^3}{(3^2)^x} = 3^{-x} \cdot 3^3 \cdot 3^{-2x} = 3^{-x+3-2x}$$

$$= 3^{3-3x} = 3^{3(1-x)}$$

7. Determine if the given function is an exponential function. If it is an exponential function, state whether it represents exponential growth or decay. $\hookrightarrow f(x) = a \cdot b^x$, $b > 0$, $b \neq 1$

(a) 5^{x+3}

$$= 5^x \cdot 5^3$$

$$= 125 \cdot 5^x$$

* exponential function

* $b = 5 > 1 \Rightarrow$ exponential growth

* if $0 < b < 1 \Rightarrow$ exp. decay
* if $b > 1 \Rightarrow$ exp. growth

(b) $-4x^{15}$

not an exponential function

* this is a polynomial / power function

(c) $7 \cdot \left(\frac{1}{3}\right)^{-2x}$

$$= 7 \cdot (3^{-1})^{-2x}$$

$$= 7 \cdot 3^{2x} = 7 \cdot (3^2)^x$$

$$= 7 \cdot 9^x$$

* exponential function

* $b = 9 > 1 \Rightarrow$ exponential growth




8. State the domain, range, end behavior, x and y -intercepts (if any) of each function below.

(a) $f(x) = \left(\frac{3}{4}\right)^{x+2}$

$$f(x) = \left(\frac{3}{4}\right)^2 \cdot \left(\frac{3}{4}\right)^x$$

$$= \frac{9}{16} \cdot \left(\frac{3}{4}\right)^x$$

* exponential decay ($b = \frac{3}{4} < 1$)
* $a = \frac{9}{16} > 0$



Domain: $(-\infty, \infty)$

Range: $(0, \infty)$

y-int: $f(0) = \frac{9}{16} \cdot \left(\frac{3}{4}\right)^0 = \frac{9}{16}$

$(0, \frac{9}{16})$

x-int: NONE


(b) $g(x) = 4^{x-2}$

$$g(x) = 4^{-2} \cdot 4^x$$

$$= \frac{1}{4} \cdot 4^x$$

$$= \frac{1}{16} \cdot 4^x$$

* exponential growth ($b = 4 > 1$)
* $a = \frac{1}{16} > 0$



Domain: $(-\infty, \infty)$

Range: $(0, \infty)$

y-int: $g(0) = \frac{1}{16} \cdot 4^0 = \frac{1}{16}$

$(0, \frac{1}{16})$

x-int: NONE



9. For each function below, state the domain using interval notation

(a) $f(x) = 3^{\frac{2x}{x+4}}$

* $f(x) = 3^{\frac{2x}{x+4}}$ is defined when $\frac{2x}{x+4}$ is defined

* $\frac{2x}{x+4}$ is a rational function \Rightarrow denom $\neq 0 \Rightarrow x+4 \neq 0 \Rightarrow x \neq -4$

Domain of f : $(-\infty, -4) \cup (-4, \infty)$

(b) $h(x) = e^{\sqrt{3-5x}}$

* $h(x) = e^{\sqrt{3-5x}}$ is defined when $\sqrt{3-5x}$ is defined

* $\sqrt{3-5x}$ is an even root function $\Rightarrow 3-5x \geq 0 \Rightarrow \frac{3}{5} \geq \frac{5x}{5} \Rightarrow \frac{3}{5} \geq x$

Domain of h : $(-\infty, \frac{3}{5}]$

(c) $g(x) = \frac{\sqrt[5]{2x-7}}{2x+3}$

* $g(x) = \frac{\sqrt[5]{2x-7}}{2x+3}$ defined when $\sqrt[5]{2x-7}$ is an odd root $\Rightarrow 2x-7$ defined $\Rightarrow x$ any real #
 $\rightarrow 2^{x+3}$ exists & $2^{x+3} \neq 0$
 \uparrow
exists for all real #s
always true since it is an exponential function
range: $(0, \infty)$

* Domain of $g(x)$: $(-\infty, \infty)$



10. Algebraically solve each equation for x

$$(a) \left(\frac{1}{8}\right)^{2x} = 16^{x-5} \Leftrightarrow (2^{-3})^{2x} = (2^4)^{x-5}$$

$$2^{-6x} = 2^{4(x-5)} = 2^{4x-20} \quad * \text{ base} = 2$$

$$-6x = 4x - 20$$

$$\frac{20}{10} = \frac{10x}{10} \Rightarrow \boxed{x = 2}$$

$$(b) 3^{x+2} = 27^{x-1} \quad 27 = 3^3$$

$$3^{x+2} = (3^3)^{x-1} = 3^{3(x-1)} = 3^{3x-3} \quad * \text{ base } 3$$

$$x+2 = 3x-3$$

$$\frac{5}{2} = \frac{2x}{2} \Rightarrow \boxed{x = \frac{5}{2}}$$

$$32 = 2^5$$

$$16 = 2^4$$

$$1 = 2^0$$

$$(c) \left(\frac{1}{32}\right) \cdot 2^{x^2} \cdot 16^x = 1$$

$$\frac{1}{2^5} \cdot 2^{x^2} \cdot 2^{4x} = 2^0 \Rightarrow 2^{-5} \cdot 2^{x^2} \cdot 2^{4x} = 2^0$$

$$2^{-5+x^2+4x} = 2^0 \quad \text{base } 2$$

$$x^2 + 4x - 5 = 0$$

$$(x+5)(x-1) = 0 \Rightarrow \boxed{x = -5, x = 1}$$



11. If you invest $\$4000$ in an account that earns interest at a rate of 3.5% per year, compounded monthly

(a) how much will be in the account after 10 years?

$$A(t) = P \left(1 + \frac{r}{m}\right)^{mt}$$

$$= 4000 \left(1 + \frac{0.035}{12}\right)^{120}$$

$$= \boxed{\$5,673.38}$$

↑ round to the nearest
CENT

- APR
- * $P = \$4000$ initial principal balance
 - * $r =$ annual interest rate $= 0.035$
 - * $m =$ # of compounding periods per year $= 12$ (monthly)
 - * $t =$ time in years $= 10$ years

(b) If the annual interest is compounded continuously instead of monthly, how much more will be in the account after 10 years compared to your previous answer?

$$A(t) = P e^{rt}$$

$$= 4000 e^{0.035 * 10}$$

$$= \boxed{\$5676.27}$$

You earn $\boxed{\$2.89}$ more, after 10 years, if interest is compounded continuously rather than monthly.



12. If $g(x) = -\frac{3}{5}f(x+5) - 9$, write the transformations that would be applied to the graph of $f(x)$ (in the correct order), to produce the graph of $g(x)$.

* Translation to the LEFT 5 units

* Vertical shrinking by a factor of $\frac{1}{(\frac{3}{5})} = \frac{5}{3}$

* Reflection across the x-axis

* Translation DOWN by 9 units

13. If the graph of $f(x) = |x|$ is shifted right 2 units, vertically expanded by a factor of 6, reflected over the x -axis, and then shifted 9 units up, what is the equation of the resulting graph?

$$f(x) = |x|$$

↓ SHIFT right 2 units

$$|x-2|$$

↓ Vertically expand by factor of 6

$$6|x-2|$$

↪ reflection over the x -axis

$$-6|x-2| \rightarrow \text{shift up by 9 units} \rightarrow -6|x-2| + 9$$

$$g(x) = -6|x-2| + 9$$