



limit as x approaches c
of $f(x)$

SESSION 1: SECTIONS 1-1 AND 1-2

y -value


The notation $\lim_{x \rightarrow c} f(x)$ (for a real number c) means we need to find the value $f(x)$ approaches when x is near, but not necessarily equal to c . The function must approach the same value (i.e., L) from both the left and right side of $x = c$ for $\lim_{x \rightarrow c} f(x) = L$

1. A graph of $f(x)$ is given below. Use the graph to find each limit below. If a limit does not exist, state so and use limit notation to describe any infinite behavior.

(a) $\lim_{x \rightarrow -4^-} f(x) = 3$ *from the left*

(b) $\lim_{x \rightarrow -4^+} f(x) = 2$ *from the right*

(c) $\lim_{x \rightarrow -4} f(x)$ DNE (Does not exist)



(d) $\lim_{x \rightarrow 2} f(x) = 5$

$\lim_{x \rightarrow 2^-} f(x) = 5$ $\lim_{x \rightarrow 2^+} f(x) = 5$

(e) $\lim_{x \rightarrow 6^+} f(x) \rightarrow -\infty$ *from the right*

(f) $\lim_{x \rightarrow 6} f(x)$ DNE

$\lim_{x \rightarrow 6^-} f(x) \rightarrow \infty$

2. Given $f(x) = \frac{5(x+3)}{x^2+5x+6}$, complete the tables below and then use the table to estimate the given limits

(a) $\lim_{x \rightarrow -3} f(x)$

| left-hand limit | | right-hand limit | |
|-----------------|--------|------------------|--------|
| x | $f(x)$ | x | $f(x)$ |
| -3.1 | -4.545 | -2.9 | -5.556 |
| -3.01 | -4.95 | -2.99 | -5.051 |
| -3.001 | -4.995 | -2.999 | -5.005 |
| -3.0001 | -5 | -2.9999 | -5.001 |

$\lim_{x \rightarrow -3^-} f(x) \approx -5$

$\lim_{x \rightarrow -3^+} f(x) \approx -5$

$\lim_{x \rightarrow -3} f(x) \approx -5$

(b) $\lim_{x \rightarrow -2} f(x)$

| left-hand limit | | right-hand limit | |
|-----------------|--------|------------------|--------|
| x | $f(x)$ | x | $f(x)$ |
| -2.1 | -50 | -1.9 | 50 |
| -2.01 | -500 | -1.99 | 500 |
| -2.001 | -5000 | -1.999 | 5000 |
| -2.0001 | -50000 | -1.9999 | 50000 |

$f(x) = \frac{5(x+3)}{x^2+5x+6}$



$\lim_{x \rightarrow -2^-} f(x) \rightarrow -\infty$

$\lim_{x \rightarrow -2^+} f(x) \rightarrow \infty$

$\lim_{x \rightarrow -2} f(x) \text{ DNE}$

Direct Substitution Property For Polynomial and Rational Functions

If P and Q are polynomials and c is any real number, then

$$\lim_{x \rightarrow c} P(x) = P(c) \quad \text{and} \quad \lim_{x \rightarrow c} \frac{P(x)}{Q(x)} = \frac{P(c)}{Q(c)}$$

as long as $Q(c)$ is nonzero.

Cases for a Ratio of Two Functions

Given two functions $f(x)$ and $g(x)$, and any real number c , use the cases below when finding $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$.

- **Case 1:** (L is any real number and $M \neq 0$)

$$\text{If } \lim_{x \rightarrow c} f(x) = L \text{ and } \lim_{x \rightarrow c} g(x) = M, \text{ then } \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}.$$

- **Case 2**

$$\text{If } \lim_{x \rightarrow c} f(x) \neq 0 \text{ and } \lim_{x \rightarrow c} g(x) = 0, \text{ then } \lim_{x \rightarrow c} \frac{f(x)}{g(x)} \text{ does not exist.}$$

- **Case 3**

If $\lim_{x \rightarrow c} f(x) = 0$ and $\lim_{x \rightarrow c} g(x) = 0$, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ cannot be determined (i.e., is indeterminate) and further algebraic manipulation is necessary to convert the limit to an expression in which Case 1 or Case 2 applies.

3. Given $h(x)$ below, find the following limits algebraically, check your results graphically.

$$h(x) = \begin{cases} 2x + 10 & x \leq -7 \\ \frac{3x - 8}{4 - x} & -7 < x < 1 \\ \frac{2x^2 + x - 3}{x^2 - 5x + 4} & x \geq 1 \end{cases}$$

left x=1

right x=1

$$(a) \lim_{x \rightarrow 1} h(x) = \boxed{\frac{-5}{3}}$$

$$\lim_{x \rightarrow 1^-} \frac{3x - 8}{4 - x} = \frac{3(1) - 8}{4 - 1} = \frac{-5}{3}$$

Because $4 - 1 = 3 \neq 0$
use direct substitution.

$$\lim_{x \rightarrow 1^+} \frac{2x^2 + x - 3}{x^2 - 5x + 4} = \lim_{x \rightarrow 1^+} \frac{(2x + 3)(\cancel{x - 1})}{(x - 4)(\cancel{x - 1})} = \lim_{x \rightarrow 1^+} \frac{2x + 3}{x - 4} = \frac{2(1) + 3}{1 - 4} = \frac{5}{-3}$$

$$\lim_{x \rightarrow 1^+} x^2 - 5x + 4 = 0$$

CASE 3

$$\lim_{x \rightarrow 1^+} 2x^2 + x - 3 = 2(1)^2 + 1 - 3 = 2 + 1 - 3 = 0$$

$$h(x) = \begin{cases} 2x + 10 & x \leq -7 \\ \frac{3x - 8}{4 - x} & -7 < x < 1 \\ \frac{2x^2 + x - 3}{x^2 - 5x + 4} & x \geq 1 \end{cases}$$

| x | $\frac{3x-8}{4-x}$ | |
|-------|--------------------|---------------|
| 3.9 | 37 | left → ∞ |
| 3.99 | 397 | |
| 3.999 | 3997 | |
| 4.1 | -43 | right → -∞ |
| 4.01 | -403 | |
| 4.001 | -4003 | |

(b) $\lim_{x \rightarrow 4} h(x) = \lim_{x \rightarrow 4} \frac{3x-8}{4-x}$ DNE

$\lim_{x \rightarrow 4} (3x-8) = 3(4) - 8 = 12 - 8 = 4$

$\lim_{x \rightarrow 4} (4-x) = 4-4 = 0$

CASE 2


but $\lim_{x \rightarrow 4^-} \frac{3x-8}{4-x} \rightarrow \infty$ and $\lim_{x \rightarrow 4^+} \frac{3x-8}{4-x} \rightarrow -\infty$

(c) $\lim_{x \rightarrow -7^-} h(x) = \lim_{x \rightarrow -7^-} (2x+10) = 2(-7) + 10 = -14 + 10 = \boxed{-4}$

(d) $\lim_{x \rightarrow 0} h(x) = \lim_{x \rightarrow 0} \frac{3x-8}{4-x} = \frac{3(0)-8}{4-0} = \boxed{\frac{-5}{4}}$

limit laws

4. Given the graph of $f(x)$ and $g(x)$ below find $\lim_{x \rightarrow -3} \left(2f(x) + \frac{g(x)}{x^2} + 8 \right)$.



$$\begin{aligned} \lim_{x \rightarrow -3} \left(2f(x) + \frac{g(x)}{x^2} + 8 \right) &= \lim_{x \rightarrow -3} 2f(x) + \lim_{x \rightarrow -3} \frac{g(x)}{x^2} + \lim_{x \rightarrow -3} 8 \\ &= 2 \lim_{x \rightarrow -3} f(x) + \frac{\lim_{x \rightarrow -3} g(x)}{(\lim_{x \rightarrow -3} x)^2} + \lim_{x \rightarrow -3} 8 \\ &= 2(4) + \frac{-5}{(-3)^2} + 8 = 8 + \frac{-5}{9} + 8 = 16 - \frac{5}{9} \\ &= 15\frac{4}{9} \end{aligned}$$

5. Find the limits below algebraically.

$$\begin{aligned} \text{(a) } \lim_{x \rightarrow -5} [\ln(6+x) - 2x] &= \ln(6+(-5)) - 2(-5) \\ &= \ln(1) + 10 \\ &= 0 + 10 \\ &= 10 \end{aligned}$$

$$\text{(b) } \lim_{x \rightarrow 4} \frac{x^2 - 8}{x + 4} = \frac{4^2 - 8}{4 + 4} = \frac{16 - 8}{8} = \frac{8}{8} = 1$$

$$(c) \lim_{x \rightarrow 4} \frac{x-4}{x+4} = \frac{4-4}{4+4} = \frac{0}{8} = 0$$

$$(d) \lim_{x \rightarrow 4} \frac{x+4}{x-4} \text{ DNE, because } \lim_{x \rightarrow 4^-} \frac{x+4}{x-4} \rightarrow -\infty, \lim_{x \rightarrow 4^+} \frac{x+4}{x-4} \rightarrow \infty.$$

$$\left. \begin{array}{l} \lim_{x \rightarrow 4} (x+4) = 4+4 = 8 \\ \lim_{x \rightarrow 4} (x-4) = 4-4 = 0 \end{array} \right\} \text{CASE 2}$$

| X | Y | |
|-------|-------|----------------|
| 3.9 | -79 | } left → -∞ |
| 3.99 | -799 | |
| 3.999 | -7999 | |
| 4.1 | 81 | } right → ∞ |
| 4.01 | 801 | |
| 4.001 | 8001 | |
| X = | | |

$$(e) \lim_{x \rightarrow 1^-} \frac{\frac{8}{x+5} - \frac{4}{x+2}}{x-1}$$

$$\lim_{x \rightarrow 1^-} \left(\frac{8}{x+5} - \frac{4}{x+2} \right) = \frac{8}{1+5} - \frac{4}{1+2} = \frac{8}{6} - \frac{4}{3} = 0$$

$$\lim_{x \rightarrow 1^-} (x-1) = 1-1=0 \quad \text{CASE 3}$$

$$\begin{aligned} \lim_{x \rightarrow 1^-} \frac{\frac{8(x+2)}{(x+5)(x+2)} - \frac{4(x+5)}{(x+2)(x+5)}}{\frac{(x-1)}{1}} &= \lim_{x \rightarrow 1^-} \left(\frac{8x+16-4x-20}{(x+5)(x+2)} \cdot \frac{1}{(x-1)} \right) \\ &= \lim_{x \rightarrow 1^-} \frac{4x-4}{(x+5)(x+2)(x-1)} \\ &= \lim_{x \rightarrow 1^-} \frac{4(x-1)}{(x+5)(x+2)(x-1)} \\ &= \lim_{x \rightarrow 1^-} \frac{4}{(x+5)(x+2)} = \frac{4}{(1+5)(1+2)} \\ &= \frac{4}{6 \cdot 3} = \frac{4}{18} = \frac{2}{9} \end{aligned}$$

$$(f) \lim_{x \rightarrow -1/2} f(x) \text{ given } f(x) = \begin{cases} \frac{2x^2 - 3x - 2}{2x + 1} & x < -\frac{1}{2} \quad \left. \vphantom{\frac{2x^2 - 3x - 2}{2x + 1}} \right\} \text{left of } x = -\frac{1}{2} \\ 2x + 7 & x > -\frac{1}{2} \quad \left. \vphantom{2x + 7} \right\} \text{right of } x = -\frac{1}{2} \end{cases}$$

$$\lim_{x \rightarrow -\frac{1}{2}^-} \frac{2x^2 - 3x - 2}{2x + 1} = \lim_{x \rightarrow -\frac{1}{2}^-} \frac{(2x+1)(x-2)}{(2x+1)} = \lim_{x \rightarrow -\frac{1}{2}^-} (x-2) = -\frac{1}{2} - 2 = -2.5$$

$$2\left(-\frac{1}{2}\right) + 1 = -1 + 1 = 0$$

$$2\left(-\frac{1}{2}\right)^2 - 3\left(-\frac{1}{2}\right) - 2 = 2\left(\frac{1}{4}\right) + \frac{3}{2} - 2$$

$$= \frac{1}{2} + \frac{3}{2} - 2$$

CASE 3

$$= 0$$

$$\lim_{x \rightarrow -\frac{1}{2}^+} (2x+7) = 2\left(-\frac{1}{2}\right) + 7 = -1 + 7 = 6$$

$$\boxed{\lim_{x \rightarrow -\frac{1}{2}} f(x) \text{ DNE}}$$